

MAGNETIC STUDIES OF AVANHANDAVA H4 AND BJURBÖLE L4 CHONDRULES

T. Kohout^{1,2,3}, G. Kletetschka^{3,4,5}, L. J. Pesonen¹ and P. J. Wasilewski⁵

1 Division of Geophysics, Department of Physical Sciences, University of Helsinki, Helsinki, Finland, tomas.kohout@helsinki.fi

2 Department of Applied Geophysics, Faculty of Science, Charles University in Prague, Prague, Czech Republic

3 Institute of Geology, Academy of Sciences of the Czech Republic, Prague, Czech Republic

4 Department of Physics, Catholic University of America, Washington DC, USA



5 Laboratory for Extraterrestrial Physics, NASA Goddard Space Flight Center, Code 691, Greenbelt, MD, USA

Introduction: The Avanhandava (H4) and Bjurböle (L4) chondrites represent primitive chondritic material of low metamorphic degree. Their friable nature allows to pick up oriented individual chondrules and to study their magnetic properties and magnetic mineralogy.

AVANHANDAVA H4

BJURBÖLE L4


Chondrule magnetic conglomerate test




The stereonet projection of the chondrule magnetic directions. The magnetic conglomerate revealed randomly magnetized chondrules and uniformly magnetized matrix.

The chondrules of the Bjurböle meteorite shows random distribution of magnetic directions.

Magnetomineralogy investigations




The magnetic carrier in Avanhandava chondrules is kamacite. Kamacite is characterized by low coercivity (<10 mT) and a-g transition at 1030 K on heating and g-a transition at 930 K on cooling curve.



Based on high coercivity of Bjurböle chondrules the tetraenaite was identified as main magnetic phase. During thermomagnetic investigations the decomposition of tetraenaite to kamacite in 750 K – 770 K temperature range was observed.

Magnetic paleofield estimate for chondrules



Paleofield estimate: The paleofield method based on the REM ratio (NRM/SIRM) [1] reveals approximate paleofields between 5 μ T and 20 μ T (REM ~ 0.002) for Avanhandava chondrules and between 12 μ T and 45 μ T (REM ~ 0.0015–0.0048) for Bjurböle chondrules. The method is based on comparison of chondrule REM ratio to the data obtained by laboratory TRM experiments with various FeNi alloys.

Conclusions: Despite the different magnetomineralogy the chondrules of both Avanhandava and Bjurböle meteorites show a low and randomly oriented NRM. The paleofields determined on chondrules are lower than geomagnetic field. That suggests together with random NRM directions that chondrules are not magnetically contaminated by geomagnetic or artificial fields and obtained their magnetization prior the conglomeration to parent body.

Acknowledgement: This work was financially supported by Academy of Finland and Väisälä Foundation at Finnish Academy of Science and Letters.



References: [1] Kletetschka G. et al, 2006; TRM in Low Magnetic Fields: a minimum field that can be recorded by large multidomain grain. Physics of the Earth and Planetary Interiors 154 (3–4), 290–298, 2006.